

BINARY BLACK HOLE MERGER: THE THEORY

INTERFACING NUMERICAL AND ANALYTICAL RELATIVITY

Alessandro Nagar

Centro Fermi, INFN, Sezione di Torino &
Institut des Hautes Etudes Scientifiques (IHES)

[nagar@ihes.fr]

The IHES effective-one-body (EOB) code: <https://eob.ihes.fr>

T. Damour, AN, S. Bernuzzi, D. Bini...

A. Nagar, 31 January 2017

GW150914

GW150914 parameters:

$$m_1 = 35.7 M_\odot$$

$$m_2 = 29.1 M_\odot$$

$$M_f = 61.8 M_\odot$$

$$a_1 \equiv S_1/(m_1^2) = 0.31^{+0.48}_{-0.28}$$


$$a_2 \equiv S_2/(m_2^2) = 0.46^{+0.48}_{-0.42}$$

$$a_f \equiv \frac{J_f}{M_f^2} = 0.67$$

$$q \equiv \frac{m_1}{m_2} = 1.27$$

Symmetric mass ratio

$$\nu \equiv \frac{m_1 m_2}{(m_1 + m_2)^2} = 0.2466$$



$$\text{strain} = \frac{\delta L}{L}$$

THE THEORY...


Is needed to compute waveform templates for characterizing the source (GWs were detected...but WHAT was detected?)

Theory is needed to study the 2-body problem in General Relativity (dynamics & gravitational wave emission)

Theory: **SYNERGY** between
Analytical and Numerical General Relativity
(AR/NR)

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G}{c^4}T_{\mu\nu}$$

UBER GRAVITATIONSWELLEN (EINSTEIN, 1918)



$$g_{ij} = \delta_{ij} + h_{ij}$$


h_{ij} is transverse and traceless and propagates at the speed of light

GRAVITATIONAL WAVES: TWO HELICITY STATES $s = \pm 2$

Massless, two helicity states,

i.e., two transverse-traceless (TT) tensor polarizations propagating at $v = c$

$$h_{ij} = h_+(x_i x_j - y_i y_j) + h_\times (x_i y_j + y_i x_j)$$



h_+

h_\times

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$$

$$\bar{h}_{\mu\nu} = h_{\mu\nu} - \frac{1}{2}\eta_{\mu\nu}h$$

$$\partial_\rho \partial^\rho \bar{h}_{\mu\nu} = -\frac{16\pi G}{c^4} T_{\mu\nu}$$

GRAVITATIONAL WAVES: PIONEERING THEIR DETECTION


Joseph Weber (1919-2000)

General Relativity and Gravitational Waves
(Interscience Publishers, NY, 1961)

$$\frac{\delta L}{L} \approx h_{ij} n^i n^j$$



LASER INTERFEROMETER GW DETECTORS



HOW TO DETECT & MEASURE: MATCHED FILTERING!

To extract/do parameter estimation of the GW signal from detector's output
(lost in broadband noise $S_n(f)$)

$$\langle output | h_{\text{template}} \rangle = \int \frac{df}{S_n(f)} o(f) h_{\text{template}}^*(f)$$



Detector's output

Template of
expected
GW signal


Need waveform templates!

GW150914


was so loud that it could be seen with the naked eye...



pass-band
filtering



OBSERVED GRAVITATIONAL WAVE SIGNALS



$$29M_{\odot} + 35M_{\odot}$$


$$23M_{\odot} + 13M_{\odot}$$

$$14M_{\odot} + 7.5M_{\odot}$$

BH radii: $\simeq 20 - 100$ km

BBHS: WAVEFORM OVERVIEW


$$h_+ - ih_\times = \frac{1}{r} \sum_{\ell m} h_{\ell m} {}_{-2}Y_{\ell m}(\theta, \phi) \quad h(m_1, m_2, \vec{S}_1, \vec{S}_2)$$




e.g: equal-mass BBH, aligned-spins

$$\chi_1 = \chi_2 = +0.98$$

- SXS (Simulating eXtreme Spacetimes) collaboration
- www.blackholes.org
- Free catalog of waveforms (downloadable)



BINARY NEUTRON STARS (BNS)?



All BNS need is Love!

$$q = 1 \quad M = 2.7M_{\odot}$$

- Tidal effects
- Love numbers (tidal “polarization” constants)
- EOS dependence & “universality”
- EOB/NR for BNS

See:

Damour&Nagar, PRD 2009

Damour&Nagar, PRD 2010

Damour,Nagar et al., PRL 2011


Bini,Damour&Faye, PRD2012

Bini&Damour, PRD 2014

Bernuzzi, Nagar, et al, PRL 2014

Bernuzzi, Nagar, Dietrich, PRL 2015


Bernuzzi, Nagar, Dietrich & Damour,PRL, 2015




FAST CHIRP: COULD GW150914 BE A BNS?

The merger occurs at frequencies too low to be a "standard" BNS

GW frequency grows from 35Hz to 150Hz around peak (factor 4) over the observed 8GWs cycles




But the final answer is that consistency was found between inspiral and ringdown!




Proposal for improved analysis with "more" ringdown: Del Pozzo&Nagar, arXiv:1606.03952

TEMPLATES FOR GWS FROM BBH COALESCENCE

Brady, Craighton & Thorne, 1998



Merger:
Numerical Relativity




Effective-One-Body (Buonanno & Damour (2000))

PN-resummation (Damour, Iyer, Sathyaprakash (1998))

Numerical Relativity: ≥ 2005 (F. Pretorius, Campanelli et al., Baker et al.)

Most accurate data: Caltech-Cornell spectral code: M. Scheel et al., 2008 (SXS collaboration)



Spectral code

Extrapolation (radius & resolution)

Phase error:



< 0.02 rad (inspiral)

< 0.1 rad (ringdown)

EFFECTIVE ONE BODY (EOB): 2000

Numerical Relativity was not working (yet...)

EOB formalism was predictive, qualitatively and semi-quantitatively correct (10%)



- Blurred transition from inspiral to plunge
- Final black-hole mass
- Final black hole spin
- Complete waveform

A. Buonanno & T. Damour, PRD 59 (1999) 084006

A. Buonanno & T. Damour, PRD 62 (2000) 064015



> 2005: Developing EOB & interfacing with NR
2 groups did (and are doing) it

- A. Buonanno et al. (AEI)
- T. Damour & AN + (>2005)

$$\nu = \frac{m_1 m_2}{(m_1 + m_2)^2} = \frac{\mu}{M}$$

PRECURSOR-BURST-RINGDOWN STRUCTURE :1972

Davis, Ruffini & Tiomno (DRT): radial plunge of a test-particle onto a Schwarzschild black hole (Regge-Wheeler-Zerilli BH perturbation theory)



NDB: TEST PARTICLE

Nagar, Damour & Bernuzzi

+Harms, Zenginoglu

RWZ: transition from inspiral to plunge

Teukolsky equation: up to very high spins

Many papers from 2006 up to today...








FIG. 1. Transition from quasicircular inspiral orbit to plunge. Initial position is $r_0 = 7M$ and $\nu = 10^{-3}$.

USEFUL AND IMPORTANT
LABORATORY TO LEARN THE
PHYSICS OF THE MERGER



IMPORTANCE OF AN ANALYTICAL FORMALISM

- **Theoretical:** physical understanding of the coalescence process, especially in complicated situations (e.g., precessing spins).
- **Practical:** need many thousands of accurate GWs templates for detection and data analysis. Need analytical templates: $h(m_1, m_2, \vec{S}_1, \vec{S}_2)$
- **Solution:** synergy between analytical & numerical relativity






BBH & BNS COALESCENCE: NUMERICAL RELATIVITY

Numerical relativity is complicated & computationally expensive:

- Formulation of Einstein equations (BSSN, harmonic, Z4c,...)
- Setting up initial data (solution of the constraints)
- Gauge choice
- Numerical approach (finite-differencing (FD, e.g. Llama) vs spectral (SpEC, SXS))
- High-order FD operators
- Treatment of BH singularity (excision vs punctures)
- Wave extraction problem on finite-size grids (Cauchy-Characteristic vs extrapolation)
- Huge computational resources (mass-ratios 1:10; spin)
- Adaptive-mesh-refinement
- Error budget (convergence rates are far from clean...)
- For BNS: further complications due to GR-Hydrodynamics for matter
- Months of running/analysis to get one accurate waveform....

Multi-patch grid structure

(Llama FD code, Pollney & Reisswig)



A catalog of 171 high-quality binary black-hole simulations for gravitational-wave astronomy [PRL 111 (2013) 241104]

Abdul H. Mroué,¹ Mark A. Scheel,² Béla Szilágyi,² Harald P. Pfeiffer,¹ Michael Boyle,³ Daniel A. Hemberger,³ Lawrence E. Kidder,³ Geoffrey Lovelace,^{4,2} Sergei Ossokine,^{1,5} Nicholas W. Taylor,² Anil Zenginoğlu,² Luisa T. Buchman,² Tony Chu,¹ Evan Foley,⁴ Matthew Giesler,⁴ Robert Owen,⁶ and Saul A. Teukolsky³




FIG. 3: Waveforms from all simulations in the catalog. Shown here are h_+ (blue) and h_x (red) in a sky direction parallel to the initial orbital plane of each simulation. All plots have the same horizontal scale, with each tick representing a time interval of $2000M$, where M is the total mass.



• www.blackholes.org

But (at least) 250.000 templates were used...

Numerical relativity reaching into post-Newtonian territory: a compact-object binary simulation spanning 350 gravitational-wave cycles

Béla Szilágyi,^{1,2} Jonathan Blackman,¹ Alessandra Buonanno,^{3,4} Andrea Taracchini,³
Harald P. Pfeiffer,^{5,6} Mark A. Scheel,¹ Tony Chu,^{7,5} Lawrence E. Kidder,⁸ and Yi Pan⁴

(Dated: February 18, 2015)



EOBNR

ANALYTICALLY: MOTION AND GW IN GR

Hamiltonian: conservative part of the dynamics

Radiation reaction: mechanical energy/angular momentum goes away in GWs and backreacts on the system.

The (closed) **orbit** **CIRCULARIZES** and **SHRINKS** with time

Waveform

General Relativity is **NONLINEAR!**

Post-Newtonian (PN) approximation: expansion in $\frac{v^2}{c^2}$

PROBLEM OF MOTION IN GENERAL RELATIVITY

Approximation methods

- ▶ post-Minkowskian (Einstein 1916)
- ▶ post-Newtonian (Droste 1916)
- ▶ Matching of asymptotic expansions: body zone/near zone/wave zone
- ▶ Numerical Relativity

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + h_{\mu\nu}(x) , \quad h_{\mu\nu} \ll 1$$
$$h_{00} \sim h_{ij} \sim \frac{v^2}{c^2} , \quad h_{0i} \sim \frac{v^3}{c^3} , \quad \partial_0 h \sim \frac{v}{c} \partial_i h$$

One-chart versus Multi-chart approaches

Coupling between Einstein field equations and equations of motion

Strongly self-gravitating bodies: neutron stars or black holes

$$h_{\mu\nu}(x) \sim 1$$

Skeletonized: $T_{\mu\nu}$ point-masses ? delta-functions in GR

Multipolar Expansion

Need to go to very high-orders of approximation

QFT-like
calculations

Use a "cocktail": PM, PN, MPM, MAE, EFT, an. reg., dim. reg.,...

POST-NEWTONIAN HAMILTONIAN (C.O.M)

$$\hat{H}_{\text{real}}^{\text{NR}}(\mathbf{q}, \mathbf{p}) = \hat{H}_{\text{N}}(\mathbf{q}, \mathbf{p}) + \hat{H}_{1\text{PN}}(\mathbf{q}, \mathbf{p}) + \hat{H}_{2\text{PN}}(\mathbf{q}, \mathbf{p}) + \hat{H}_{3\text{PN}}(\mathbf{q}, \mathbf{p}), \quad (4.27)$$

where

$$\hat{H}_{\text{N}}(\mathbf{q}, \mathbf{p}) = \frac{\mathbf{p}^2}{2} - \frac{1}{q}, \quad \text{Newton (OPN)} \quad (4.28a)$$

$$\hat{H}_{1\text{PN}}(\mathbf{q}, \mathbf{p}) = \frac{1}{8}(3\nu - 1)(\mathbf{p}^2)^2 - \frac{1}{2}[(3 + \nu)\mathbf{p}^2 + \nu(\mathbf{n} \cdot \mathbf{p})^2] \frac{1}{q} + \frac{1}{2q^2}, \quad (1\text{PN}, 1938) \quad (4.28b)$$

- [Einstein-Infeld-Hoffman]

$$\begin{aligned} \hat{H}_{2\text{PN}}(\mathbf{q}, \mathbf{p}) = & \frac{1}{16}(1 - 5\nu + 5\nu^2)(\mathbf{p}^2)^3 + \frac{1}{8}[(5 - 20\nu - 3\nu^2)(\mathbf{p}^2)^2 - 2\nu^2(\mathbf{n} \cdot \mathbf{p})^2\mathbf{p}^2 - 3\nu^2(\mathbf{n} \cdot \mathbf{p})^4] \frac{1}{q} \\ & + \frac{1}{2}[(5 + 8\nu)\mathbf{p}^2 + 3\nu(\mathbf{n} \cdot \mathbf{p})^2] \frac{1}{q^2} - \frac{1}{4}(1 + 3\nu)\frac{1}{q^3}, \quad (2\text{PN}, 1982/83) \quad (4.28c) \end{aligned}$$

- [Damour-Deruelle]

$$\begin{aligned} \hat{H}_{3\text{PN}}(\mathbf{q}, \mathbf{p}) = & \frac{1}{128}(-5 + 35\nu - 70\nu^2 + 35\nu^3)(\mathbf{p}^2)^4 \\ & + \frac{1}{16}[(-7 + 42\nu - 53\nu^2 - 5\nu^3)(\mathbf{p}^2)^3 + (2 - 3\nu)\nu^2(\mathbf{n} \cdot \mathbf{p})^2(\mathbf{p}^2)^2 + 3(1 - \nu)\nu^2(\mathbf{n} \cdot \mathbf{p})^4\mathbf{p}^2 - 5\nu^3(\mathbf{n} \cdot \mathbf{p})^6] \frac{1}{q} \\ & + \left[\frac{1}{16}(-27 + 136\nu + 109\nu^2)(\mathbf{p}^2)^2 + \frac{1}{16}(17 + 30\nu)\nu(\mathbf{n} \cdot \mathbf{p})^2\mathbf{p}^2 + \frac{1}{12}(5 + 43\nu)\nu(\mathbf{n} \cdot \mathbf{p})^4 \right] \frac{1}{q^2} \quad (3\text{PN}, 2000) \\ & + \left\{ \left[-\frac{25}{8} + \left(\frac{1}{64}\pi^2 - \frac{335}{48} \right) \nu - \frac{23}{8}\nu^2 \right] \mathbf{p}^2 + \left(-\frac{85}{16} - \frac{3}{64}\pi^2 - \frac{7}{4}\nu \right) \nu(\mathbf{n} \cdot \mathbf{p})^2 \right\} \frac{1}{q^3} \\ & + \left[\frac{1}{8} + \left(\frac{109}{12} - \frac{21}{32}\pi^2 + \omega_{\text{static}} \right) \nu \right] \frac{1}{q^4}. \quad (4.28d) \end{aligned}$$

- [Damour, Jaranowski, Schaefer]

...and **4PN** too, [Damour, Jaranowski&Schaefer 2014/2015] - 4 loop calculation

$$\mathbf{q} = \mathbf{q}_1 - \mathbf{q}_2$$

$$\mathbf{p} = \mathbf{p}_1 = -\mathbf{p}_2$$

PN-EXPANDED (CIRCULAR) ENERGY FLUX (3.5PN)

$$\frac{dE}{dt} = -\mathcal{L}$$

balance equation

Mechanical loss

GW luminosity

$$\begin{aligned} \mathcal{L} = \frac{32c^5}{5G} \nu^2 x^5 \left\{ 1 + \left(-\frac{1247}{336} - \frac{35}{12}\nu \right) x + 4\pi x^{3/2} + \left(-\frac{44711}{9072} + \frac{9271}{504}\nu + \frac{65}{18}\nu^2 \right) x^2 \right. \\ \text{Newtonian} \\ \text{quadrupole} \\ \text{formula} \\ \left. + \left(-\frac{8191}{672} - \frac{583}{24}\nu \right) \pi x^{5/2} \right. \\ + \left[\frac{6643739519}{69854400} + \frac{16}{3}\pi^2 - \frac{1712}{105}C - \frac{856}{105}\ln(16x) \right. \\ \left. + \left(-\frac{134543}{7776} + \frac{41}{48}\pi^2 \right) \nu - \frac{94403}{3024}\nu^2 - \frac{775}{324}\nu^3 \right] x^3 \\ \left. + \left(-\frac{16285}{504} + \frac{214745}{1728}\nu + \frac{193385}{3024}\nu^2 \right) \pi x^{7/2} + \mathcal{O}\left(\frac{1}{c^8}\right) \right\}. \end{aligned}$$

$$C = \gamma_E = 0.5772156649\dots$$

TAYLOR-EXPANDED (CIRCULAR) 3PN WAVEFORM

Blanchet, Iyer&Joguet, 02; Blanchet, Damour, Iyer&Esposito-Farese, 04; Kidder07; Blanchet et al.,08

$$\begin{aligned}
 h^{22} = & -8\sqrt{\frac{\pi}{5}} \frac{G\nu m}{c^2 R} e^{-2i\phi} x \left\{ 1 - x \left(\frac{107}{42} - \frac{55}{42} \nu \right) + x^{3/2} \left[2\pi + 6i \ln\left(\frac{x}{x_0}\right) \right] - x^2 \left(\frac{2173}{1512} + \frac{1069}{216} \nu - \frac{2047}{1512} \nu^2 \right) \right. \\
 & - x^{5/2} \left[\left(\frac{107}{21} - \frac{34}{21} \nu \right) \pi + 24i\nu + \left(\frac{107i}{7} - \frac{34i}{7} \nu \right) \ln\left(\frac{x}{x_0}\right) \right] \\
 & + x^3 \left[\frac{27\,027\,409}{646\,800} - \frac{856}{105} \gamma_E + \frac{2}{3} \pi^2 - \frac{1712}{105} \ln 2 - \frac{428}{105} \ln x \right. \\
 & \left. \left. - 18 \left[\ln\left(\frac{x}{x_0}\right) \right]^2 - \left(\frac{278\,185}{33\,264} - \frac{41}{96} \pi^2 \right) \nu - \frac{20\,261}{2772} \nu^2 + \frac{114\,635}{99\,792} \nu^3 + \frac{428i}{105} \pi + 12i\pi \ln\left(\frac{x}{x_0}\right) \right] + O(\epsilon^{7/2}) \right\},
 \end{aligned}$$

$$x = (M\Omega)^{2/3} \sim v^2/c^2$$

$$M = m_1 + m_2$$

$$\nu = \frac{m_1 m_2}{M^2}$$

EFFECTIVE-ONE-BODY (EOB)

approach to the general relativistic two-body problem

(Buonanno-Damour 99, 00, Damour-Jaranowski-Schäfer 00, Damour 01, Damour-Nagar 07, Damour-Iyer-Nagar 08)


key ideas:

- (1) Replace two-body dynamics (m_1, m_2) by dynamics of a particle ($\mu \equiv m_1 m_2 / (m_1 + m_2)$) in an effective metric $g_{\mu\nu}^{\text{eff}}(u)$, with

$$u \equiv GM/c^2 R, \quad M \equiv m_1 + m_2$$

- (2) Systematically use **RESUMMATION** of PN expressions (both $g_{\mu\nu}^{\text{eff}}$ and \mathcal{F}_{RR}) based on various physical requirements
- (3) Require **continuous deformation w.r.t.**
 $\nu \equiv \mu/M \equiv m_1 m_2 / (m_1 + m_2)^2$ in the interval $0 \leq \nu \leq \frac{1}{4}$

STRUCTURE OF THE EOB FORMALISM



TWO-BODY/EOB “CORRESPONDENCE”:

THINK QUANTUM-MECHANICALLY (J.A. WHEELER)


Real 2-body system
(in the c.o.m. frame)
(m_1, m_2)



An effective particle
in some effective metric

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$

$$g_{\mu\nu}^{\text{eff}}$$



$$\mu^2 + g_{\text{eff}}^{\mu\nu} \frac{\partial S_{\text{eff}}}{\partial x^\mu} \frac{\partial S_{\text{eff}}}{\partial x^\nu} + \mathcal{O}(p^4) = 0$$

Sommerfeld's
“Old Quantum Mechanics”
(action-angle variables &
Delaunay Hamiltonian)

$$J = \ell \hbar = \frac{1}{2\pi} \oint p_\varphi d\varphi$$

$$N = n \hbar = I_r + J$$

$$I_r = \frac{1}{2\pi} \oint p_r dr$$

$$H^{\text{classical}}(q, p) \longrightarrow H^{\text{classical}}(I_a) \longrightarrow E^{\text{quantum}}(I_a = n_a h) = f^{-1}[\mathcal{E}_{\text{eff}}^{\text{quantum}}(I_a^{\text{eff}} = n_a h)]$$

THE EOB ENERGY MAP

$$\mu = \frac{m_1 m_2}{m_1 + m_2}$$

Real 2-body system
(PN-expanded Hamiltonian
in the c.o.m. frame)

(m_1, m_2)

1:1 map

An effective particle
in some effective metric

$g_{\mu\nu}^{\text{eff}}$

Simple energy map:

$$\mathcal{E}_{\text{eff}} = \frac{E_{\text{real}}^2 - m_1^2 - m_2^2}{2M}$$

EOB Hamiltonian:

$$H_{\text{EOB}} = M \sqrt{1 + 2\nu \left(\hat{H}_{\text{eff}} - 1 \right)}$$

$$M = m_1 + m_2$$

$$\nu = \frac{\mu}{M}$$

$$\hat{H}_{\text{eff}} = \frac{H_{\text{eff}}}{\mu}$$

EXPLICIT FORM OF THE EOB HAMILTONIAN

EOB Hamiltonian

$$H_{\text{EOB}} = M \sqrt{1 + 2\nu \left(\hat{H}_{\text{eff}} - 1 \right)}$$

All functions are a ν -dependent deformation of the Schwarzschild ones

$$A(r) = 1 - 2u + 2\nu u^3 + a_4 \nu u^4$$

$$a_4 = \frac{94}{3} - \frac{41}{32} \pi^2 \simeq 18.6879027$$

$$A(r)B(r) = 1 - 6\nu u^2 + 2(3\nu - 26)\nu u^3 \quad u = GM/(c^2 R)$$

Simple effective Hamiltonian:

$$\hat{H}_{\text{eff}} \equiv \sqrt{p_{r_*}^2 + A(r) \left(1 + \frac{p_\varphi^2}{r^2} + z_3 \frac{p_{r_*}^4}{r^2} \right)} \quad p_{r_*} = \left(\frac{A}{B} \right)^{1/2} p_r$$

Crucial EOB radial potential

Contribution at 3PN

EFFECTIVE POTENTIALS

Newtonian gravity (any mass ratio):
circular orbits are always stable. No plunge.

$$W_{\text{Newt}}^{\text{eff}} = 1 - \frac{2}{r} + \frac{p_{\varphi}^2}{r^2}$$

Test-body on Schwarzschild black hole:



last stable orbit (LSO) at $r=6M$; plunge

$$W_{\text{Schwarzschild}}^{\text{eff}} = \left(1 - \frac{2}{r}\right) \left(1 + \frac{p_{\varphi}^2}{r^2}\right)$$

EOB, Black-hole binary, any mass ratio:

last stable orbit (LSO) at $r < 6M$ plunge


$$W_{\text{EOB}}^{\text{eff}} = A(r; \nu) \left(1 + \frac{p_{\varphi}^2}{r^2}\right)$$



ν -deformation of the Schwarzschild case!

HAMILTON'S EQUATIONS & RADIATION REACTION

$$\begin{aligned}\dot{r} &= \left(\frac{A}{B}\right)^{1/2} \frac{\partial \hat{H}_{\text{EOB}}}{\partial p_{r_*}} \\ \dot{\varphi} &= \frac{\partial \hat{H}_{\text{EOB}}}{\partial p_{\varphi}} \equiv \Omega \\ \dot{p}_{r_*} &= - \left(\frac{A}{B}\right)^{1/2} \frac{\partial \hat{H}_{\text{EOB}}}{\partial r} + \hat{\mathcal{F}}_{r_*} \\ \dot{p}_{\varphi} &= \hat{\mathcal{F}}_{\varphi}\end{aligned}$$




- The system must radiate angular momentum
- How? Use PN-based (Taylor-expanded) radiation reaction force (ang-mom flux)
- Need flux resummation

$$\hat{\mathcal{F}}_{\varphi}^{\text{Taylor}} = -\frac{32}{5} \nu \Omega^5 r_{\Omega}^4 \hat{F}^{\text{Taylor}}(v_{\varphi}) \rightarrow$$

Plus horizon contribution [AN&Akcaay2012]

Resummation multipole by multipole
(Damour&Nagar 2007,
Damour, Iyer & Nagar 2008,
Damour & Nagar, 2009)

USE OF PADE APPROXIMANTS



- Continuity with Schwarzschild metric: $A(r)$ needs to have a zero
- Simple (possible) prescription: use a Padé representation of the potential

$$A(r) = P_3^1[A^{3\text{PN}}(r)] = \frac{1 + n_1 u}{1 + d_1 u + d_2 u^2 + d_3 u^3}$$

MULTIPOLAR WAVEFORM RESUMMATION

Resummation of the waveform (and flux) multipole by multipole (**CRUCIAL!**)

[Damour&Nagar 2007, Damour, Iyer, Nagar 2008]

Next-to-quasi-circular correction

$$h_{\ell m} \equiv \underbrace{h_{\ell m}^{(N, \epsilon)}}_{\text{Newtonian}} \underbrace{\hat{h}_{\ell m}^{(\epsilon)}}_{\text{PN-correction}} \underbrace{\hat{h}_{\ell m}^{\text{NQC}}}_{\text{NQC}}$$

Newtonian \times PN \times NQC

$$\hat{h}_{\ell m}^{(\epsilon)} = \hat{S}_{\text{eff}}^{(\epsilon)} T_{\ell m} e^{i\delta_{\ell m}} \rho_{\ell m}^{\ell}$$

Remnant phase and modulus corrections: "improved" PN series

Effective source:

EOB (effective) energy (even-parity modes)

EOB angular momentum (odd-parity modes)



The "Tail factor"

$$T_{\ell m} = \frac{\Gamma(\ell + 1 - 2i\hat{k})}{\Gamma(\ell + 1)} e^{\pi\hat{k}} e^{2i\hat{k} \ln(2kr_0)}$$

Resums an infinite number of leading logarithms in tail effects (hereditary contributions)


EFFECTIVENESS OF FLUX RESUMMATION

Test-mass
(Comparing fluxes,
circular orbits)



Equal-mass
(Comparing non-resummed & EOB-
resummed amplitudes to Caltech-Cornell
BBH data)

$$\mathcal{F}_\varphi \equiv -\frac{1}{8\pi\Omega} \sum_{\ell=2}^{\ell_{\max}} \sum_{m=1}^{\ell} (m\Omega)^2 |Rh_{\ell m}^{(\epsilon)}|^2$$



THE KNOWLEDGE OF THE CENTRAL A POTENTIAL TODAY

4PN analytically complete + 5PN logarithmic term in the $A(u)$ function:

[Damour 2009, Blanchet et al. 2010, Barack, Damour & Sago 2010, Le Tiec et al. 2011, Barausse et al. 2011, Akcay et al. 2012, Bini & Damour 2013, Damour Jaranowski & Schaefer 2014].

$$A_{5\text{PN}}^{\text{Taylor}} = \underbrace{1}_{1\text{PN}} - \underbrace{2u}_{2\text{PN}} + \underbrace{2\nu u^3}_{3\text{PN}} + \underbrace{\left(\frac{94}{3} - \frac{41}{32}\pi^2\right)\nu u^4}_{4\text{PN}} + \underbrace{\nu[a_5^c(\nu) + a_5^{\ln} \ln u]u^5}_{4\text{PN}} + \underbrace{\nu[a_6^c(\nu) + a_6^{\ln} \ln u]u^6}_{5\text{PN}}$$

$$\left. \begin{aligned} a_5^{\log} &= \frac{64}{5} \\ a_5^c &= a_{50}^c + \nu a_{51}^c \\ a_{50}^c &= -\frac{4237}{60} + \frac{2275}{512}\pi^2 + \frac{256}{5}\log(2) + \frac{128}{5}\gamma \\ a_{51}^c &= -\frac{221}{6} + \frac{41}{32}\pi^2 \end{aligned} \right\} \text{4PN fully known ANALYTICALLY!}$$


$$a_6^{\log} = -\frac{7004}{105} - \frac{144}{5}\nu \quad \text{5PN logarithmic term (analytically known)}$$

NEED ONE "effective" 5PN parameter from NR waveform data: $a_6^c(\nu)$

State-of-the-art EOB potential (5PN-resummed):

$$A(u; \nu, a_6^c) = P_5^1[A_{5\text{PN}}^{\text{Taylor}}(u; \nu, a_6^c)]$$

THE EOB[NR] POTENTIAL




From EOB/NR-fitting: $a_6^c(\nu) = 3097.3\nu^2 - 1330.6\nu + 81.3804$

TAKE AWAY:

BBH system is more bound, smaller "separation" and higher frequencies!

NDRP, arXiv:1506.08457

RESULTS: EOBNR/NR WAVEFORMS (NO SPIN)



equal-mass case

Nagar, Damour, Reisswig & Pollney, PRD 93 (2016), 04404

HIGHER MODES (NO SPIN)

- Unpublished, but free to download at eob.ihes.fr (Matlab code)
- Check unfaithfulness vs NR surrogate
(G. Pratten & AN, 2016 in preparation)





FIG. 4. Unfaithfulness for the EOB waveforms against surrogate waveforms as a function of the total mass of the system M for mass ratios $q = 2, 4, 6, 8$ and 10 assuming $\theta = \phi = \pi/3$. The top plot shows a comparison of multimodal waveforms constructed from (22), (21) and (33). The bottom plot shows a comparison for waveforms constructed from just the (22) modes.




SPINNING BBHs

Spin-orbit & spin-spin couplings

(i) Spins **aligned** with **L**: **repulsive** (slower) **L-o-n-g-e-r INSPIRAL**

(ii) Spins **anti-aligned** with **L**: **attractive** (faster) **shorter INSPIRAL**

(iii) **Misaligned spins**: precession of the orbital plane (**waveform modulation**)



$$\chi_{1,2} = \frac{c \mathbf{S}_{1,2}}{G m_{1,2}^2}$$


EOB/NR agreement: sophisticated (though rather simple) model for spin-aligned binaries

Damour&Nagar, PRD90 (2014), 024054 (Hamiltonian)

Damour&Nagar, PRD90 (2014), 044018 (Ringdown)

Nagar,Damour, Reisswig & Pollney, PRD 93 (2016), 044046

AEI model, SEOBNRv4, Bohe et al., arXiv:1611.03703v1 (PRD in press)



SO & SS EFFECTS IN EOB HAMILTONIAN

New way of combining available knowledge within some Hamiltonian

[Damour&Nagar, PRD 2014]

$$\hat{H}_{\text{eff}} = \frac{g_S^{\text{eff}}}{r^3} \mathbf{L} \cdot \mathbf{S} + \frac{g_{S^*}^{\text{eff}}}{r^3} \mathbf{L} \cdot \mathbf{S}^* + \sqrt{A(1 + \gamma^{ij} p_i p_j + Q_4(p))}$$

with the structure

$$g_S^{\text{eff}} = 2 + \nu(\text{PN corrections}) + (\text{spin})^2 \text{corrections}$$

$$g_{S^*}^{\text{eff}} = \left(\frac{3}{2} + \text{test mass coupling} \right) + \nu(\text{PN corrections}) + (\text{spin})^2 \text{corrections}$$

$$A = 1 - \frac{2}{r} + \nu(\text{PN corrections}) + (\text{spin})^2 \text{corrections}$$

$$\gamma^{ij} = \gamma_{\text{Kerr}}^{ij} + \nu(\text{PN corrections}) + \dots$$

$$\mathbf{S} = \mathbf{S}_1 + \mathbf{S}_2 = M^2 (X_1^2 \chi_1 + X_2^2 \chi_2) \quad X_i = m_i/M$$

$$\mathbf{S}^* = \frac{m_2}{m_1} \mathbf{S}_1 + \frac{m_1}{m_2} \mathbf{S}_2 = M^2 \nu (\chi_1 + \chi_2) \quad -1 \leq \chi_i \leq 1$$

THE TWO TYPES OF SPIN-ORBIT COUPLINGS

$$\hat{H}_{\text{SO}}^{\text{eff}} = G_S \mathbf{L} \cdot \mathbf{S} + G_{S^*} \mathbf{L} \cdot \mathbf{S}^* \quad G_S = \frac{1}{r^3} g_S^{\text{eff}}, \quad G_{S^*} = \frac{1}{r^3} g_{S^*}^{\text{eff}}$$

In the Kerr limit, only **S-type gyro-gravitomagnetic ratio** enters:

$$g_S^{\text{eff}} = 2 \frac{r^2}{r^2 + a^2 \left[(1 - \cos^2 \theta) \left(1 + \frac{2}{r} \right) + 2 \cos^2 \theta \right] + \frac{a^4}{r^2} \cos^2 \theta} = 2 + \mathcal{O}[(\text{spin})^2]$$

PN calculations yield (in some spin gauge)[DJS08, Hartung&Steinhoff11, Nagar11, Barausse&Buonanno11]

$$\begin{aligned} g_S^{\text{eff}} &= 2 + \frac{1}{c^2} \left\{ -\frac{15}{r} \nu - \frac{33}{8} (\mathbf{n} \cdot \mathbf{p})^2 \right\} \\ &\quad + \frac{1}{c^4} \left\{ -\frac{1}{r^2} \left(\frac{51}{4} \nu + \frac{\nu^2}{8} \right) + \frac{1}{r} \left(-\frac{21}{2} \nu + \frac{23}{8} \nu^2 \right) (\mathbf{n} \cdot \mathbf{p})^2 + \frac{5}{8} \nu (1 + 7\nu) (\mathbf{n} \cdot \mathbf{p})^4 \right\}, \quad + \frac{1}{c^6} \frac{\nu c_3}{r^3} \\ g_{S^*}^{\text{eff}} &= \frac{3}{2} + \frac{1}{c^2} \left\{ -\frac{1}{r} \left(\frac{9}{8} + \frac{3}{4} \nu \right) - \left(\frac{9}{4} \nu + \frac{15}{8} \right) (\mathbf{n} \cdot \mathbf{p})^2 \right\} \\ &\quad + \frac{1}{c^4} \left\{ -\frac{1}{r^2} \left(\frac{27}{16} + \frac{39}{4} \nu + \frac{3}{16} \nu^2 \right) + \frac{1}{r} \left(\frac{69}{16} - \frac{9}{4} \nu + \frac{57}{16} \nu^2 \right) (\mathbf{n} \cdot \mathbf{p})^2 + \left(\frac{35}{16} + \frac{5}{2} \nu + \frac{45}{16} \nu^2 \right) (\mathbf{n} \cdot \mathbf{p})^4 \right\} + \frac{1}{c^6} \frac{\nu c_3}{r^3} \end{aligned}$$

"Effective" NNNLO SO-coupling

The NR-informed effective parameter makes the spin-orbit coupling stronger or weaker with respect to the simple analytical prediction

40 NR SXS Datasets (public in the fall of 2013 and used before for SEOBNRv2)

TABLE I: EOB/NR phasing comparison. The columns report: the number of the dataset; the name of the configuration in the SXS catalog; the mass ratio $q = m_1/m_2$; the symmetric mass ratio ν ; the dimensionless spins χ_1 and χ_2 ; the phase difference $\Delta\phi^{\text{EOBNR}} \equiv \phi^{\text{EOB}} - \phi^{\text{NR}}$ computed at NR merger; the NR phase uncertainty at NR merger $\delta\phi_{\text{mrg}}^{\text{NR}}$ (when available) measured taking the difference between the two highest resolution levels (see text); the maximum value of the unfaithfulness $\bar{F} \equiv 1 - F$ as per Eq. (22). The $\Delta\phi^{\text{EOBNR}}$'s in brackets for $\chi_1 = \chi_2 > +0.85$ were obtained using Eq. (21) for $\Delta t^{\text{NQC}}(\chi)$.

#	Name	N orbits	q	ν	χ_1	χ_2	$\Delta\phi_{\text{mrg}}^{\text{EOBNR}}$ [rad]	$\delta\phi_{\text{mrg}}^{\text{NR}}$ [rad]	$\max(\bar{F})$
1	SXS:BBH:none	14	1	0.25	0.0	0.0	-0.016	...	0.00087
2	SXS:BBH:0066	28	1	0.25	0.0	0.0	+0.010	...	0.00068
3	SXS:BBH:0002	32.42	1	0.25	0.0	0.0	+0.073	0.066	0.00101
4	SXS:BBH:0007	29.09	1.5	0.24	0	0	+0.05	0.018	0.00201
5	SXS:BBH:0169	15.68	2	$0.\bar{2}$	0	0	-0.15	0.02	0.00045
6	SXS:BBH:0030	18.22	3	0.1875	0	0	-0.074	0.087	0.00035
7	SXS:BBH:0167	15.59	4	0.16	0	0	-0.059	0.52	0.00035
8	SXS:BBH:0056	28.81	5	0.13 $\bar{8}$	0	0	-0.089	0.44	0.00038
9	SXS:BBH:0166	21.56	6	0.1224	0	0	-0.198	...	0.00037
10	SXS:BBH:0063	25.83	8	0.0987	0	0	-0.453	1.01	0.00292
11	SXS:BBH:0185	24.91	9.98911	0.0827	0	0	-0.0051	0.376	0.00066
12	SXS:BBH:0004	30.19	1	0.25	-0.50	0.0	-0.017	0.068	0.00403
13	SXS:BBH:0005	30.19	1	0.25	+0.50	0.0	+0.08	0.28	0.00052
14	SXS:BBH:0156	12.42	1	0.25	-0.95	-0.95	+0.32	2.17	0.00058
15	SXS:BBH:0159	12.67	1	0.25	-0.90	-0.90	+0.06	0.38	0.00047
16	SXS:BBH:0154	13.24	1	0.25	-0.80	-0.80	+0.11	...	0.00044
17	SXS:BBH:0151	14.48	1	0.25	-0.60	-0.60	-0.049	0.14	0.00042
18	SXS:BBH:0148	15.49	1	0.25	-0.44	-0.44	+0.14	0.72	0.00043
19	SXS:BBH:0149	17.12	1	0.25	-0.20	-0.20	+0.45	0.90	0.00085
20	SXS:BBH:0150	19.82	1	0.25	+0.20	+0.20	+0.94	0.99	0.00275
21	SXS:BBH:0152	22.64	1	0.25	+0.60	+0.60	+0.01	0.36	0.00068
22	SXS:BBH:0155	24.09	1	0.25	+0.80	+0.80	-0.39	0.26	0.00110
23	SXS:BBH:0153	24.49	1	0.25	+0.85	+0.85	+0.06	...	0.00059
24	SXS:BBH:0160	24.83	1	0.25	+0.90	+0.90	+0.41 (+0.41)	0.80	0.00117
25	SXS:BBH:0157	25.15	1	0.25	+0.95	+0.95	+0.37 (+0.83)	1.18	0.00295
26	SXS:BBH:0158	25.27	1	0.25	+0.97	+0.97	+0.37 (+0.49)	1.26	0.00325
27	SXS:BBH:0172	25.35	1	0.25	+0.98	+0.98	+0.99 (+0.46)	2.02	0.00422
28	SXS:BBH:0177	25.40	1	0.25	+0.99	+0.99	+0.22 (+0.48)	0.40	0.00507
29	SXS:BBH:0178	25.43	1	0.25	+0.994	+0.994	+0.24 (+0.23)	-0.53	0.00506
30	SXS:BBH:0013	23.75	1.5	0.24	+0.5	0	+0.31	...	0.00058
31	SXS:BBH:0014	22.63	1.5	0.24	-0.5	0	-0.15	0.15	0.00046
32	SXS:BBH:0162	18.61	2	$0.\bar{2}$	+0.6	0	-0.20	0.71	0.00027
33	SXS:BBH:0036	31.72	3	0.1875	-0.5	0	+0.08	0.065	0.00040
34	SXS:BBH:0031	21.89	3	0.1875	+0.5	0	+0.12	0.034	0.00023
35	SXS:BBH:0047	22.72	3	0.1875	+0.5	+0.5	-0.034	...	0.00030
36	SXS:BBH:0046	14.39	3	0.1875	-0.5	-0.5	+0.36	...	0.00054
37	SXS:BBH:0110	24.24	5	0.13 $\bar{8}$	+0.5	0	+0.24	...	0.00016
38	SXS:BBH:0060	23.17	5	0.13 $\bar{8}$	-0.5	0	+0.21	0.8	0.00034
39	SXS:BBH:0064	19.16	8	0.0987	-0.5	0	+0.026	0.8	0.00042
40	SXS:BBH:0065	33.97	8	0.0987	+0.5	0	+1.33	-3.0	0.00040


Several equal-mass,
equal-spin data

Just a few unequal-
mass, unequal-spin data

SPIN-ORBIT NR INFORMATION

Procedure:

- (i) align waveforms in the early inspiral;
- (ii) tune the parameter to have phase difference compatible with the NR uncertainty




+ interpolating fits for NQC functioning point, ringdown coefficients etc.
(Achille's heel...still ok..)

$$\tilde{a}_{1,2} = X_{1,2} \chi_{1,2}$$

$$X_{1,2} \equiv \frac{m_{1,2}}{M}$$

Quasi-linear function of the spins

$$c_3(\tilde{a}_1, \tilde{a}_2, \nu) = p_0 \frac{1 + n_1(\tilde{a}_1 + \tilde{a}_2) + n_2(\tilde{a}_1 + \tilde{a}_2)^2}{1 + d_1(\tilde{a}_1 + \tilde{a}_2)} + (p_1\nu + p_2\nu^2 + p_2\nu^3)(\tilde{a}_1 + \tilde{a}_2)\sqrt{1 - 4\nu} + p_4(\tilde{a}_1 - \tilde{a}_2)\nu^2,$$





EOBNR MODEL USED FOR GW150914

Different EOB Hamiltonian [Barausse & Buonanno11, Taracchini et al.12]

SEOBNRv2: Taracchini, Buonanno et al., PRD 89, 061502 (R), 2014

SEOBNRv2_ROM_DoubleSpin: M. Puerrer, CQG 31, 195010 (2014)



Effectively used to get the masses:

SEOBNRv2_ROM_DoubleSpin


IMRPhenom (Khan et al., 2015)

just AFTER, the best choices
were cross checked with NR simulations!


IHES EOBNR MODEL

Best existing EOBNR model **WAS NOT** used for parameter estimation:
EOB/EOBNR UNFAITHFULNESS (40 NR SXS dataset)

SEOBNRv2



IHSEOspin






$$\bar{F} \equiv 1 - \max_{t_0, \phi_0} \frac{\langle h_{22}^{\text{EOB}}, h_{22}^{\text{NR}} \rangle}{||h_{22}^{\text{EOB}}|| ||h_{22}^{\text{NR}}||}$$

$$\langle h_1, h_2 \rangle \equiv 4\Re \int_{f_{\min}}^{\infty} \tilde{h}_1(f) \tilde{h}_2^*(f) / S_n(f) df$$


Nagar, Damour, Reisswig & Pollney, PRD 93 (2016), 044046

FIRST QUESTION: MEASURING PARAMETERS




ROBUSTNESS?

SEOBNRv2



SEOBNRv4



grey: below 3%

AEI model: Bohe et al. arXiv: 1611.03703v1

4 parameters

Strong recalibration of the state-of-the-art SEOBNRv2 model (used for O1) to have it faithful towards a set of 141 NR simulations (about 100 new ones)

More NR simulations seem essential to "calibrate & improve" the AEI EOBNR model


$$d_{\text{SO}} = +147.481449\chi^3v^2 - 568.651115\chi^3v + 66.198703\chi^3 - 343.313058\chi^2v + 2495.293427\chi v^2 - 44.532373,$$

$$d_{\text{SS}} = +528.511252\chi^3v^2 - 41.000256\chi^3v + 1161.780126\chi^2v^3 - 326.324859\chi^2v^2 + 37.196389\chi v + 706.958312v^3 - 36.027203v + 6.068071,$$

BUT THIS IS NOT GENERAL...

October 31st: 93 NR datasets released publicly. These are those used to calibrate SEOBNRv4 (+ others non public)
First use them to cross-check our model.


Interpolating NR fits for NQC point & ringdown. Previous NR data plus (5,-0.90,0)



Our EOBNR model is very robust and consistent
ALSO outside the "information" domain over
93 new waveforms. Three outliers above
1% (but always below 3%).

3% Better performance than SEOBNRv2 with
no need of further NR information

1%




MINIMAL RECALIBRATION

Best value of the c_3 parameter for the three outliers. Check phase agreement in the time-domain to be within the NR error bar.
New fit to the best values to determine new values of the parameters of the unequal-mass sector.


Recalibration with 3 more NR datasets; 90 datasets as a cross/check.

Done by hand, no need of sophisticated mechanisms/algorithms. **IMPROVABLE: NQC & RINGDOWN FITS USING MORE NR DATA**



Nagar, Riemenschneider & Pratten, 2017, in prep.

WHAT TO IMPROVE?



More NR data sets to be included both in the NQC-functioning-point fit as well as in the postmerger fit (see Del Pozzo & Nagar, arXiv:1606.03952). This is an easily solvable problem (in progress).

It is reasonable to aim at 0.1% level unfaithfulness. This is easily at reach of the model.
More precise "calibration" and/or improved theoretical structures.

PRECESSION

Different EOB Hamiltonian [Barausse & Buonanno11, Taracchini et al.12]

SEOBNRv3: Taracchini, Buonanno et al., PRD 89, 061502 (R), 2014

Babak, Taracchini & Buonanno, 2016




FIG. 9: We show for cases 3 and 4 of Table I the GW polarization h_+ , containing contributions from $\ell = 2$ modes, that propagates along a direction \hat{N} specified by spherical coordinates $\theta = \pi/3$ and $\phi = \pi/2$ associated with the inertial source frame $\{e_1^S, e_2^S, e_3^S\}$. The EOB waveforms start at the after-junk-radiation times of $t = 230M$ and $t = 160M$, respectively.

Good EOBNR/NR agreement.

The method works

Slow: analysis is time-consuming

Improvements in the implementation are needed

PhenomP: P. Schmidt et al. 2012/2014

Phenomenological Precessing model that takes into account precession effects at leading order by “twisting” nonprecessing waveforms.

Conclusion: no precession could be really seen.

POSTMERGER DESCRIPTION



Damour&AN, PRD 2014: motivated because the "standard" QNMs attachment is far from trivial for high-spins

Originally conceived for EOB; useful also as a stand-alone postmerger template

Del Pozzo & AN, arXiv: 1606.03952

ANALYTIC TEMPLATE for the FULL POSTMERGER signal coming from a suitable fit of NR data.

$$\sigma_1 = \alpha_1 + i\omega_1$$



EFFECTIVE FIT

Damour&AN 2014

Factorize the fundamental

QNM, fit what remains

$$h(\tau) = e^{\sigma_1 \tau - i\phi_0} \bar{h}(\tau)$$

$$\bar{h}(\tau) \equiv A_{\bar{h}} e^{i\phi_{\bar{h}}(\tau)}.$$

$$A_{\bar{h}}(\tau) = c_1^A \tanh(c_2^A \tau + c_3^A) + c_4^A,$$

$$\phi_{\bar{h}}(\tau) = -c_1^\phi \ln \left(\frac{1 + c_3^\phi e^{-c_2^\phi \tau} + c_4^\phi e^{-2c_2^\phi \tau}}{1 + c_3^\phi + c_4^\phi} \right)$$

$$c_2^A = \frac{1}{2} \alpha_{21},$$

$$\alpha_{21} = \alpha_2 - \alpha_1$$



$$c_4^A = \hat{A}_{22}^{\text{mrg}} - c_1^A \tanh(c_3^A),$$

$$c_1^A = \hat{A}_{22}^{\text{mrg}} \alpha_1 \frac{\cosh^2(c_3^A)}{c_2^A},$$

$$c_1^\phi = \Delta\omega \frac{1 + c_3^\phi + c_4^\phi}{c_2^\phi (c_3^\phi + 2c_4^\phi)},$$

$$\Delta\omega \equiv \omega_1 - M_{\text{BH}} \omega_{22}^{\text{mrg}}$$

$$c_2^\phi = \alpha_{21},$$





Good performance of primary fits (modulo details...)

Do this for various SXS dataset and then build up a (simple-minded) interpolating fit



Black-list:

- (1) the structure due to $m < 0$ modes is not included (yet)
- (2) large-mass ratios/high spin: amplitude problems
- (3) problems are extreme for high-spin EMRL waves
- (4) more flexible fit-template needed
- (5) improve/check over all datasets (SXS & BAM for large mass-ratios & consistency with EMRL)

TESTS



Phase alignment@mrg



Time&phase shift alignment (as template)

WAVEFORM RECONSTRUCTION





FIG. 4: From top to bottom right panels: GE case reconstructed post-merger waveform and corresponding 90% confidence region for SXS:BBH:0305 with post-merger SNR = 10, 20, 50 and 100. On the left hand side CO reconstructed post-merger waveform and corresponding 90% confidence region for SXS:BBH:0305 with post-merger SNR = 10, 20, 50 and 100. In all cases, the post-merger waveform is reconstructed very accurately, with uncertainty decreasing as the post-merger SNR increases.






MEASURING

GW150914-like signal

No priors on
individual masses



Priors on
individual masses




QNMS

TABLE II: Dataset of the SXS catalog used for the cross-validation of the template waveform, see Fig. 3. The last two columns list fundamental QNMs frequency inferred from NR data and measured with the post-merger template, after adding to the NR waveform some Gaussian noise. For all waveforms, we fixed the post-merger SNR = 10. The uncertainty on the measured quantities corresponds to the 90% credible regions. The datasets marked with an * were used in the construction of the template

ID	q	ν	$S_1/(m_1)^2$	$S_2/(m_2)^2$	M_{BH}/M	$J_{\text{BH}}/M_{\text{BH}}^2$	σ_1^{NR}	$\sigma_1^{\text{measured}}$
*	1	0.25	0	0	0.95161	0.6864	$0.0813 + i0.527$	$0.07_{-0.01}^{+0.02} + i0.52_{-0.06}^{+0.06}$
SXS:BBH:0152*	1	0.25	+0.60	+0.60	0.9269	0.8578	$0.0706 + i0.629$	$0.06_{-0.02}^{+0.02} + i0.64_{-0.07}^{+0.06}$
SXS:BBH:0211	1	0.25	+0.90	-0.90	0.9511	0.6835	$0.081 + i0.525$	$0.06_{-0.02}^{+0.02} + i0.50_{-0.06}^{+0.05}$
SXS:BBH:0178*	1	0.25	+0.994	+0.994	0.8867	0.9499	$0.053 + i0.746$	$0.08_{-0.02}^{+0.03} + i0.74_{-0.07}^{+0.08}$
SXS:BBH:0305	1.221	0.2475	+0.3300	-0.4399	0.9520	0.6921	$0.081 + i0.529$	$0.07_{-0.03}^{+0.05} + i0.55_{-0.06}^{+0.06}$
SXS:BBH:0025	1.5	0.2400	+0.4995	-0.4995	0.9504	0.7384	$0.079 + i0.550$	$0.08_{-0.03}^{+0.04} + i0.56_{-0.07}^{+0.06}$
SXS:BBH:0184	2	$0.\bar{2}$	0	0	0.9612	0.6234	$0.083 + i0.502$	$0.28_{-0.22}^{+0.20} + i0.53_{-0.39}^{+0.41}$
SXS:BBH:0162	2	$0.\bar{2}$	+0.6000	0	0.9461	0.8082	$0.075 + i0.591$	$0.08_{-0.03}^{+0.04} + i0.56_{-0.07}^{+0.08}$
SXS:BBH:0257	2	$0.\bar{2}$	+0.85	+0.85	0.9199	0.9175	$0.062 + i0.694$	$0.07_{-0.02}^{+0.03} + i0.67_{-0.08}^{+0.07}$
SXS:BBH:0045	3	0.1875	+0.4995	-0.4995	0.9628	0.7410	$0.079 + i0.552$	$0.21_{-0.18}^{+0.26} + i0.59_{-0.45}^{+0.36}$
SXS:BBH:0292	3	0.1875	+0.7314	-0.8493	0.9560	0.8266	$0.073 + i0.604$	$0.08_{-0.02}^{+0.03} + i0.58_{-0.07}^{+0.07}$
SXS:BBH:0293	3	0.1875	+0.85	+0.85	0.9142	0.9362	$0.062 + i0.689$	$0.07_{-0.02}^{+0.03} + i0.67_{-0.07}^{+0.07}$
SXS:BBH:0317	3.327	0.1777	0.5226	-0.4482	0.9642	0.7462	$0.078 + i0.554$	$0.06_{-0.02}^{+0.02} + i0.55_{-0.06}^{+0.05}$
SXS:BBH:0208*	5	$0.13\bar{8}$	-0.90	0	0.98822	-0.12817	$0.089 + i0.359$	$0.11_{-0.02}^{+0.02} + i0.40_{-0.04}^{+0.04}$
SXS:BBH:0203	7	0.1094	+0.40	0	0.9836	0.6056	$0.083 + i0.495$	$0.07_{-0.01}^{+0.02} + i0.48_{-0.04}^{+0.06}$
SXS:BBH:0207	7	0.1094	-0.60	0	0.9909	-0.0769	$0.089 + i0.364$	$0.08_{-0.01}^{+0.02} + i0.35_{-0.04}^{+0.04}$
SXS:BBH:0064*	8	0.0987	-0.50	0	0.9922	-0.0526	$0.089 + i0.367$	$0.09_{-0.05}^{+0.12} + i0.46_{-0.08}^{+0.11}$
SXS:BBH:0185	9.990	0.0827	0	0	0.9917	0.2608	$0.087 + i0.412$	$0.12_{-0.03}^{+0.04} + i0.42_{-0.06}^{+0.07}$

BINARY NEUTRON STARS (BNS)?



All BNS need is Love!

$$q = 1 \quad M = 2.7M_{\odot}$$

- Tidal effects
- Love numbers (tidal “polarization” constants)
- EOS dependence & “universality”
- EOB/NR for BNS

See:

Damour&Nagar, PRD 2009

Damour&Nagar, PRD 2010

Damour,Nagar et al., PRL 2011


Bini,Damour&Faye, PRD2012

Bini&Damour, PRD 2014

Bernuzzi, Nagar, et al, PRL 2014

Bernuzzi, Nagar, Dietrich, PRL 2015

Bernuzzi, Nagar, Dietrich & Damour,PRL, 2015



MEASURING LOVE NUMBERS



<2012. Inspiral only; not very promising [Hinderer et al. + 2008]

IMPORTANT RESULT (Damour ,Nagar, Villain 2012)

Tidal polarizability parameters
can actually be measured by
adv LIGO with a reasonable
SNR=16

Use EOB controlled, accurate,
description of the phasing
up to BNS merger!

Confermed by Bayesian analysis:
Del Pozzo+ 2013 Agathos+2015



A. Nagar - 18 May 2016

THREE RESULTS

1. **Numerical-relativity** matches effective-one-body (EOB) **analytical-relativity** waveforms and dynamics essentially up to merger. Method to compute GW templates for LIGO/Virgo to measure EOS out of tidal effects

S. Bernuzzi, A. Nagar, T. Dietrich & T. Damour, PRL 114 (2015), 161103

"Modeling the Dynamics of Tidally Interacting Binary Neutron Stars up to Merger"

[checked by Hotokezaka et al., PRD 91 (2015) 6, 064060, notably with reduced eccentricity]

2. **Quasi-universality** in BNS merger (binding energy, angular momentum, GW frequency vs tidal coupling constant): explained using EOB theory

S. Bernuzzi, A. Nagar, S. Balmelli, T. Dietrich & M. Ujevic, PRL 112 (2014), 201101

"Quasiuniversal properties of neutron star mergers"

3. **Quasi-universality** of post-merger Mf_2 frequency vs tidal coupling constant

S. Bernuzzi, A. Nagar & T. Dietrich, arXiv:1504.01764

"Towards a description of the complete gravitational wave spectrum of neutron star mergers"

Unifying description of inspiral, merger and post-merger phases

BNS:ANALYTICAL NEEDS

- Study the response of each neutron star to the tidal field of the companion [theory of **relativistic Love numbers** (i.e. tidal polarizability coefficients) + **tidal corrections to dynamics** (beyond Newtonian accuracy)]
- Incorporate the corresponding tidal effects within a theoretical framework able to describe the gravitational wave signal emitted by inspiralling compact binaries (**possibly up to merger**): **EOB-resummed** description of dynamics and waveforms
- Compare analytical models against NR simulations, possibly calibrating high-order tidal corrections if needed
- Assess the **measurability of tidal effects** within the signal seen by interferometric detectors

LOVE NUMBERS IN GENERAL RELATIVITY

Relativistic star in an external **gravito-electric** & **gravito-magnetic (multipolar)** tidal field



The star acquires induced gravito-electric and gravito-magnetic multipole moments

Linear tidal polarization

**Induced
multipole
moments**

$$M_L^{(A)}$$

$$=$$

$$\mu_\ell^A G_L^{(A)}$$

$$S_L^{(A)}$$

$$=$$

$$\sigma_\ell^A H_L^{(A)}$$

**External
multipolar
field**

$$G\mu_\ell$$

$$=$$

$$[length]^{2\ell+1}$$

$$G\sigma_\ell$$

$$=$$

$$[length]^{2\ell+1}$$

$$2k_\ell \equiv (2\ell - 1)!! \frac{G\mu_\ell}{R^{2\ell+1}}$$

$$j_\ell \equiv (2\ell - 1)!! \frac{4(\ell + 2)}{\ell - 1} \frac{G\sigma_\ell}{R^{2\ell+1}}$$

Dimensionless relativistic
"second" Love numbers

Actual calculation based on star perturbation theory: Love numbers are obtained as boundary conditions (matching interior to exterior perturbations)

A. Nagar - 18 May 2016 - GGI

RELATIVISTIC LOVE NUMBERS (POLYTROPIC EOS)

"rest-mass polytrope" (solid lines)

$$p = K\mu^\gamma$$

$$e = \mu + \frac{p}{\gamma - 1}$$

"energy polytrope" (dashed lines)

$$p = Ke^\gamma$$


Tidal polarization parameters

$$M_L^{(A)} = \mu_\ell^A G_L^{(A)}$$

$$2k_\ell \equiv (2\ell - 1)!! \frac{G\mu_\ell}{R^{2\ell+1}}$$

Newtonian
values

Newtonian
values



EOBNR
65


Relativistic
values

RELATIVISTIC LOVE NUMBERS: (REALISTIC EOS)

DN 2009 (SLy & FPS)

Hinderer et al. 2010 (several tabulated EOS)

Polytropic EOS



$$p = K \rho^{1+1/n}$$

$$p = K \epsilon^{1+1/n}$$

A. Nagar - 06 May 2014 - Torino

TIDAL EFFECTS IN EOB FORMALISM

Tidal extension of EOB formalism: **nonminimal worldline couplings**

$$\Delta S_{\text{nonminimal}} = \sum_A \frac{1}{4} \mu_2^A \int ds_A (u^\mu u^\nu R_{\mu\alpha\nu\beta})^2 + \dots$$

Damour&Esposito-Farèse96, Goldberger&Rothstein06, TD&AN09

Relativistic
Love number

Modifications of the EOB effective metric...

$$\begin{aligned} A(r) &= A_r^0 + A^{\text{tidal}}(r) \\ \underline{A^{\text{tidal}}(r)} &= -\kappa_2^T u^6 (1 + \bar{\alpha}_1 u + \bar{\alpha}_2 u^2 + \dots) + \dots \end{aligned}$$

And tidal modifications of GW waveform & radiation reaction


- Need analytical theory for computing $\mu_2, \kappa_2^T, \bar{\alpha}_1 \dots$
- (?) Need accurate NR simulations to “calibrate” the higher-order PN tidal contributions, that may be quite important during the late inspiral

TIDAL INTERACTION POTENTIAL

Tidal “coupling constant”:

$$\kappa_\ell^T \equiv 2 \left[\frac{1}{q} \left(\frac{X_A}{C_A} \right)^{2\ell+1} k_\ell^A + q \left(\frac{X_B}{C_B} \right)^{2\ell+1} k_\ell^B \right]$$

$$X_{A,B} \equiv M_{A,B}/M$$



$$\begin{array}{l} M_A = M_B \\ \ell = 2 \\ k_2^A = k_2^B \end{array} \quad \kappa_2^T = \frac{1}{8} \frac{k_2}{C^5}$$

Function of: masses, compactnesses and relativistic Love numbers

In the dynamics:

$$A(u) = A^0(u) + A^{\text{tidal}}$$

$$A^{\text{tidal}} = \sum_{\ell \geq 2} -\kappa_\ell^T u^{2\ell+2} \hat{A}_\ell^{\text{tidal}}(u)$$

“Newtonian” (LO) part
+ PN corrections (NLO, NNLO, ...)

$$\kappa_2^T \sim 100$$

NLO & NNLO tidal PN corrections known analytically

[Bini, Damour & Faye 2011]

$$\hat{A}_2^{\text{tidal}} = 1 + \frac{5}{4}u + \frac{85}{14}u^2$$


RESUMMED TIDAL INTERACTION

Bini&Damour (2015) resummed expression for $\hat{A}_\ell^{\text{tidal}}$

Presence of a pole: potential strongly attractive @ mrg

$$A_T^{(+)}(u; \nu) \equiv - \sum_{\ell=2}^4 \left[\kappa_A^{(\ell)} u^{2\ell+2} \hat{A}_A^{(\ell+)} + (A \leftrightarrow B) \right]$$

$$\hat{A}_A^{(2+)}(u) = 1 + \frac{3u^2}{1 - r_{\text{LR}}u} + \frac{X_A \tilde{A}_1^{(2+)} \text{1SF}}{(1 - r_{\text{LR}}u)^{7/2}} + \frac{X_A^2 \tilde{A}_2^{(2+)} \text{2SF}}{(1 - r_{\text{LR}}u)^p}$$






FIG. 2: Energetics: comparison between NR data, $\text{TEOB}_{\text{Resum}}$, $\text{TEOB}_{\text{NNLO}}$ and TPN. Each bottom panel shows the two EOB-NR differences. The filled circles locate the merger points (top) and the corresponding differences (bottom). The shaded area indicates the NR uncertainty. The $\text{TEOB}_{\text{Resum}}$ model displays, globally, the smallest discrepancy with NR data (notably for merger quantities), supporting the theoretical, light-ring driven, amplification of the relativistic tidal factor.

S. Bernuzzi, A. Nagar, T. Dietrich & T. Damour, PRL 114 (2015), 161103

A. Nagar - 18 May 2016 - GGI

Waveform






FIG. 3: Phasing and amplitude comparison (versus NR retarded time) between $\text{TEOB}_{\text{Resum}}$, NR and the phasing of TT4 for three representative models. Waves are aligned on a time window (vertical dot-dashed lines) corresponding to $I_\omega \approx (0.04, 0.06)$. The markers in the bottom panels indicate: the crossing of the $\text{TEOB}_{\text{Resum}}$ LSO radius; NR (also with a dashed vertical line) and EOB merger moments.

Name	EOS	κ_2^T	τ_{LR}	$\mathcal{C}_{A,B}$	$M_{A,B}[M_\odot]$	$M_{\text{ADM}}^0[M_\odot]$	$\mathcal{J}_{\text{ADM}}^0[M_\odot^2]$	$\Delta\phi_{\text{NRmrg}}^{\text{TT4}}$	$\Delta\phi_{\text{NRmrg}}^{\text{TEOB}_{\text{NNLO}}}$	$\Delta\phi_{\text{NRmrg}}^{\text{TEOB}_{\text{Resum}}}$	$\delta\phi_{\text{NRmrg}}^{\text{NR}}$
2B135	2B	23.9121	3.253	0.2049	1.34997	2.67762	7.66256	-1.25	-0.19	+0.57 ^a	± 4.20
SLy135	SLy	73.5450	3.701	0.17381	1.35000	2.67760	7.65780	-2.75	-1.79	-0.75	± 0.40
Γ_2164	$\Gamma = 2$	75.0671	3.728	0.15999	1.64388	3.25902	11.11313	-2.29	-1.36	-0.31	± 0.90
Γ_2151	$\Gamma = 2$	183.3911	4.160	0.13999	1.51484	3.00497	9.71561	-2.60	-1.92	-1.27	± 1.20
H4135	H4	210.5866	4.211	0.14710	1.35003	2.67768	7.66315	-3.02	-2.43	-1.88	± 1.04
MS1b135	MS1b	289.8034	4.381	0.14218	1.35001	2.67769	7.66517	-3.25	-2.84	-2.45	± 3.01

EOS_{QUASI}-UNIVERSALITY OF BNS MERGER

S. Bernuzzi, A. Nagar, S. Balmelli, T. Dietrich and M. Ujevic, PRL 112 (2014) 201101

- GW frequency
- binding energy
- angular momentum
- GW amplitude
-



κ_2^T

A. Nagar - Creta 2015

EOS QUASI-UNIVERSALITY OF BNS MERGER

Matter effects on binary neutron star waveforms

Jocelyn S. Read,^{1,2} Luca Baiotti,^{3,4} Jolien D. E. Creighton,⁵ John L. Friedman,⁵ Bruno Giacomazzo,⁶ Koutarou Kyutoku,⁵ Charalampos Markakis,^{7,9} Luciano Rezzolla,⁸ Masaru Shibata,⁴ and Keisuke Taniguchi¹⁰





FIG. 4. Instantaneous gravitational-wave frequency at the point of peak amplitude, as a function of the tidal parameter $\Lambda^{1/5}$ (bottom panel) and as a function of individual star compactness C (top). For each model, the highest-resolution simulation for a given EOS is plotted in black, lower-resolution simulations in grey. The $x = (\pi M f)^{2/3} = C$ relation used in [15] to characterize merger frequency is shown in the compactness plot. An empirical fit using $\Lambda^{1/5}$ is shown in the bottom plot; the frequency of merger is more tightly correlated with Λ than with compactness/radius.



$$\Lambda = \frac{2}{3} k_2 \frac{1}{C^5}$$

empirical fit using $\Lambda^{1/5}$????

A. Nagar - Creta 2015

EOS QUASI-UNIVERSALITY OF BNS MERGER

Simple EOS-universal behavior:
measure the frequency, constrain the EOS




$$\kappa_\ell^T \equiv 2 \left[\frac{1}{q} \left(\frac{X_A}{C_A} \right)^{2\ell+1} k_\ell^A + q \left(\frac{X_B}{C_B} \right)^{2\ell+1} k_\ell^B \right]$$

$$M\Omega = \dot{\Phi} = \partial_{p_\varphi} H_{\text{EOB}}(p, q; \kappa_T)$$

A. Nagar - Creta 2015

EOS _{QUASI}-UNIVERSALITY OF BNS MERGER

Putting spins & changing the mass ratio






Test of principle:

Using 4PN, purely analytical, tidal EOB model.

Qualitatively OK; Quantitatively, differences due to the simplified dynamics

A. Nagar - Creta 2015

NEW T-EOB MODEL



Good quantitative
agreement

$$M\omega_{22}^{\text{mrg}} \approx 2\Omega_{\text{mrg}} = \left. \frac{dE_b}{dj} \right|_{j=j_{\text{mrg}}}$$

- Creta 2015

INSPIRAL-MERGER-POSTMERGER??




TABLE I: BNS configurations and data. Columns: Configuration name, equation of state, binary mass M , mass ratio q , f_2 frequency in kHz, tidal coupling constant κ_2^T .

Name	EOS	$M [M_\odot]$	q	f_2 [kHz]	κ_2^T
SLy-135135	SLy	2.70	1.00	3.48	74
SLy-145125	SLy	2.70	1.16	3.42	75
ENG-135135	ENG	2.70	1.00	2.86	91
SLy-140120	SLy	2.60	1.17	3.05	96
MPA1-135135	MPA1	2.70	1.00	2.57	115
SLy-140110	SLy	2.50	1.27	2.79	126
ALF2-135135	ALF2	2.70	1.00	2.73	138
ALF2-145125	ALF2	2.70	1.16	2.66	140
H4-135135	H4	2.70	1.00	2.50	211
H4-145125	H4	2.70	1.16	2.36	212
ALF2-140110	ALF2	2.50	1.27	2.38	216
MS1b-135135	MS1b	2.70	1.00	2.00	290
MS1-135135	MS1	2.70	1.00	1.95	327
MS1-145125	MS1	2.70	1.16	2.06	331
MS1b-140110	MS1b	2.50	1.27	2.08	441
2H-135135	2H	2.70	1.00	1.87	439
MS1b-150100	MS1b	2.50	1.50	1.87	461

S. Bernuzzi, A. Nagar & T. Dietrich, arXiv:1504.01764

OUTLOOK

1. NR/EOB (IMRPhenom is also EOB based) is the way to go. NON resummed templates are useless. Same for BNS up to merger
2. **EOB_IHES_spin**: Analytical freedom: **only two flexibility parameters** that are extracted from NR data as simple (separate) functions of symmetric mass ratio and spin magnitude
3. Compatibility (within NR errors) between such EOBNR model and state-of-the art NR data over mass ratio and spin (+precession using SEOBNRv3 exists)
4. Improvements needed: best templates, **were NOT used for analyses** (though this is irrelevant now). **This will be done in the next future on the Virgo/INFN side**


THE 2-BODY HAMILTONIAN [2PN]

The 2-body Hamiltonian at 2PN (c.o.m. frame)

$$H_{2\text{PN}}^{\text{relative}}(\mathbf{q}, \mathbf{p}) = H_0(\mathbf{q}, \mathbf{p}) + \frac{1}{c^2} H_2(\mathbf{q}, \mathbf{p}) + \frac{1}{c^4} H_4(\mathbf{q}, \mathbf{p})$$

The Newtonian limit

$$H_0(\mathbf{q}, \mathbf{p}) = \frac{\mathbf{p}^2}{2\mu} + \frac{GM\mu}{|\mathbf{q}|}$$



4 additional terms at 1PN
7 additional terms at 2PN
11 additional terms at 3PN

Rewrite the c.o.m. (reduced, non-relativistic) energy using action variables

Obtain the 2PN "quantum" energy levels: Delaunay Hamiltonian [Damour-Schaefer 1988]

$$E_{2\text{PN}}^{\text{NR}} = -\frac{1}{2}\mu \frac{\alpha^2}{n^2} \left[1 + \frac{\alpha^2}{c^2} \left(\frac{c_{11}}{n\ell} + \frac{c_{20}}{n^2} \right) + \frac{\alpha^4}{c^4} \left(\frac{c_{13}}{n\ell^3} + \frac{c_{22}}{n^2\ell^2} + \frac{c_{31}}{n^3\ell} + \frac{c_{40}}{n^4} \right) \right]$$

Balmer formula!

$$\alpha = (GM\mu)/\hbar$$

$$N = n\hbar$$

$$e \equiv \mu \quad E_n = -\frac{\mu}{2} \frac{e^4}{\hbar^2 n^2}$$

$$Ze \equiv GM$$

$$E_{2\text{PN}}^{\text{relativistic}}(n, \ell) = Mc^2 + E_{2\text{PN}}^{\text{NR}}(n, \ell)$$