



## **Brian Punsly**

Position: Research Scientist

Period covered: 6/2012-6/2013:

### **I Scientific Work**

#### Black Holes and Quasars

##### 1. Introduction

This report describes the research performed by Brian Punsly and collaborators in cooperation with ICRANet in 2012-2013. There were two lines of research. The first was directed at finding environmental factors that are related to the switch-on of the general relativistic engine responsible for a few percent of quasars driving powerful relativistic jets. This is important since this will related directly to constraints on the initial state and boundary conditions on numerical models of black hole driven jets.. The second area of research is based on using the jet in the Galactic black hole, GRS 1915+105, as a test case for black hole driven jets.

##### 2. AGN Environments and the Launching of Jets

In 2012, the research was concentrated on the nature of the broad emission line gas in AGN that launch relativistic jets. I am also leading collaborations to perform high frequency (high resolution), time resolved VLBA observations of broad absorption line quasars. Broad absorption line quasars have weak or no central engine for powerful radio jets with the jets rarely strong enough to make it out of the host galaxy. As principal investigator, in collaboration with Paola Marziani and Giovanna Stirpe at Istituto Nazionale di Astrofisica and Shaohua Zhang, we were granted telescope time on the VLT of the European Southern Observatory to study the H $\beta$  line widths of quasars with broad high ionization absorption line flows for the first time. This is a follow-up to previous work with Shaohua Zhang on quasars with low ionization broad absorption lines that indicated narrow H $\beta$  line widths consistent with polar outflows (Punsly and Zhang 2010)..

##### 2a. Asymmetric Line profiles in Blazars

The Astrophysical Journal Letter, describes the paradoxical occurrence of redward asymmetric broad emission lines in blazars which have relativistic jets point nearly pole-on towards earth. Hence these are the most blueshifted objects in the known Universe. They also have the most redshifted known wings in their broad emission lines. This odd occurrence of redshifted structures within a highly blueshifted system provides valuable clues to the origin of relativistic jets in some quasars.

Abstract: Multi-Epoch Observations of the Redwing Excess in the Spectrum of 3C279:

It has been previously determined that there is a highly significant correlation between the spectral index from 10 GHz to 1350 Å and the amount of excess luminosity in the red wing of quasar CIV λ1549 broad emission lines (BELs). Ostensibly, the prominence of the red excess is associated with the radio jet emission mechanism and is most pronounced for lines of sight close to the jet axis. Studying the scant significant differences in the UV spectra of radio loud and radio quiet quasars might provide vital clues to the origin of the unknown process that creates powerful relativistic jets that appear in only a few percent of quasars. In this study, the phenomenon is explored with multi-epoch observations of the MgII λ2798 broad line in 3C 279 which has one of the largest known redwing excesses in a quasar spectrum. The amount of excess that is detected appears to be independent of all directly observed optical continuum, radio or submm properties (fluxes or polarizations). The only trend that occurs in this sparse data is: the stronger the BEL, the larger the fraction of flux that resides in the redwing. It is concluded that more monitoring is needed and spectropolarimetry with a large telescope is essential during low states to understand more.

2b. VLBA Observations of Sub-Parsec Structure in Mrk 231: Interaction between a Relativistic Jet and a BAL Wind

I am leading an effort to study Mrk 231 at the highest resolution. It is the nearest broad absorption line quasar and we have proven that it conforms with the idea of a polar broad absorption line outflow (instead of the popular notion of an equatorial outflow) that was developed in Punsly (1999a,b). This research and proposal is being done in collaboration with Cormac Reynolds (Curtin University of Technology, Department of Imaging and Applied Physics), Christopher P. O'Dea (Department of Physics, Rochester Institute of Technology) and Joan Wrobel (NRAO, Socorro).

2b.1. Large VLBA Proposal Approved

We have been approved annually for the past few years for a very aggressive observation this object.

Abstract

We propose VLBA monitoring at 8.4, 15, 22 and 43 GHz of a high frequency flare in the nearby quasar MRK231. The "target of opportunity" observation (ToO) would be triggered by a flare detected by VLA monitoring at 22 and 43 GHz (see related proposal). The primary goals would be to detect a superluminal motion, estimate the internal energy of the flare from the spectrum and component sizes, and monitor the temporal evolution in order to understand the energy injection mechanism (rise) and the cooling mechanism (decay).

## Background

From previous VLBA studies of MRK231 in Reynolds et al (2009) and other RQ (radio quiet) quasar studies, we have seen that RQ AGN can have relativistic outflows with significant kinetic luminosities (but maybe for short periods of time). So this raises the question what is it that makes some sources RQ and others radio loud (RL)? At a redshift of 0.042, MRK231 is one of the nearest radio quiet quasars to earth. The radio core is perhaps the brightest of any radio quiet quasar at high frequency (22 and 43 GHz). The combination of significant 43 GHz flux density and its proximity to earth makes MRK231 the optimal radio quiet quasar for study with VLBA. No other radio quiet quasar central engine can be explored with such high resolution, so it is ideal for studying the high kinetic luminosity relativistic ejecta in radio quiet quasars. 43 GHz VLBA observations can fully resolve nuclear structure to within  $3.5 \times 10^{17}$  cm. We propose to use sensitive high resolution observations to study the temporal evolution of the size and spectrum of a strong flare in MRK231 in order to shed light on why such strong flares cool off and never link to large scale powerful radio lobes.

### 2b2. VLBA Observations of Parsec Scale Structure of the “Radio Loud” BALQSO FIRST J1556+3517

I am also leading an effort to study FIRST J1556+3517 at the high resolution. It is one of the nearest broad absorption line quasar and we have proven (Ghosh and Punshly 2007) that it conforms with the idea of a polar broad absorption line outflow (instead of the popular notion of an equatorial outflow) that was developed in Punshly (1999a,b). The first epoch observations are complete the second epoch observations are still in the proposal review cycle.. This proposal was done in collaboration with Cormac Reynolds (Curtin University of Technology, Department of Imaging and Applied Physics), and Christopher P. O'Dea (Department of Physics, Rochester Institute of Technology).

ABSTRACT FROM ACCEPTED PROPOSAL: We propose VLBA observations at 1.8, 5, 8.4 and 15 GHz of the Broad Absorption Line Quasar FIRST J1556+3517 (“the first radio loud BALQSO”). The primary goal would be to resolve the flat spectrum radio core for the first time. Determination of the radio jet direction, in consort with the knowledge that the jet is relativistic and viewed in a pole-on orientation and the known PA of the optical continuum polarization tightly restrict the quasar geometry. This will allow us to directly constrain the relative orientations of the “dusty torus” (scattering surface), accretion disk and the broad absorption line outflow. We also propose multiple frequency observations to look for free-free absorption that might arise from the local environment of the accretion disk or the BAL wind gas itself. If the jet is resolved by the VLBA, this observation would be the first data point in a search for component motion. If the jet is not resolved, the incredibly compact nature of the relativistic outflow indicates a severe kinematical environment.

### 3. GRS 1915+105 as a Laboratory for Studying Black Hole Driven Jets

I am currently embarked on a research program to study the Galactic black hole jet in GRS 1915+105. There is much confusion in this field because it is led by scientist not familiar with the history of astrophysical jets or the theory of black holes. There are two large projects that were

developed to understand the relationship of the energy output to the state of the accretion flow when the jets are launched. The first paper was published early this year, Punsly and Rodriguez (2013), with collaborator Jerome Rodriguez of Laboratoire AIM, CEA/DSM-CNRS-Universit\{e} Paris Diderot, IRFU SAp, F-91191 Gif-sur-Yvette, France.

ABSTRACT from The Relationship Between X-ray Luminosity and Major Flare Launching in GRS~1915+105:

We perform the most detailed analysis to date of the X-ray state of the Galactic black hole candidate GRS~1915+105 just prior to (0 to 4 hours) and during the brief (1 to 7 hour) ejection of major (superluminal) radio flares. A very strong model independent correlation is found between the 1.2 keV - 12 keV X-ray flux 0 to 4 hours before flare ejections with the peak optically thin 2.3 GHz emission of the flares. This suggests a direct physical connection between the energy in the ejection and the luminosity of the accretion flow preceding the ejection. In order to quantify this concept, we develop techniques to estimate the intrinsic (unabsorbed) X-ray luminosity,  $L_{\text{intrinsic}}$ , from RXTE ASM data and to implement known methods to estimate the time averaged power required to launch the radio emitting plasmoids,  $Q$  (sometimes called jet power). We find that the distribution of intrinsic luminosity from 1.2 keV - 50 keV,  $L_{\text{intrinsic}} (1.2 - 50)$ , is systematically elevated just before ejections compared to arbitrary times when there are no major ejections. The estimated  $Q$  is strongly correlated with  $L_{\text{intrinsic}} (1.2 - 50)$ , 0 to 4 hours before the ejection, the increase in  $L_{\text{intrinsic}} (1.2 - 50)$ , in the hours preceding the ejection and the time averaged  $L_{\text{intrinsic}} (1.2 - 50)$ , during the flare rise. Furthermore, the total time averaged power during the ejection ( $Q$  + the time average of  $L_{\text{intrinsic}} (1.2 - 50)$ , during ejection) is strongly correlated with  $L_{\text{intrinsic}} (1.2 - 50)$ , just before launch with near equality if the distance to the source is  $\approx 10$  kpc.

The second paper accepted in ApJ this year looks at the results above in the context of quasars and 3-D numerical simulations of black holes.

ABSTRACT FROM: GRS 1915+105 as a Galactic Analog of a Fanaroff-Riley II Quasar

We study the long term time averaged kinetic luminosity,  $Q$ , of the major flares of the Galactic microquasar GRS 1915+105 and the relationship to the intrinsic X-ray (bolometric) luminosity,  $L_{\text{bol}}$ , and scale it to that of a complete sample of SDSS/FIRST FR II quasars. If the scale invariance hypothesis for black holes (BHs) holds then we show that the expected distribution in the  $Q - L_{\text{bol}}$  scatter plane of GRS 1915+105 is consistent with FR II quasars for distances  $D = 10.7 - 11$  kpc. We compare the specific values of kinetic luminosity and  $L_{\text{bol}}$  during flares of GRS 1915+105 to that predicted by several 3-D MHD simulations of BH accretion flows with relativistic ejections. If FR II quasars are a scaled up version of GRS 1915+105, the data are consistent with numerical models when they contain an ergospheric disk jet and the BH spin is  $a/M = 0.99$  or  $a/M = 0.998$  (we estimate  $a/M > 0.984$ ). In the framework of scale invariance of BHs, our results may imply that FR II quasars also hold rapidly rotating BHs.

### **2013 List of Publication**

Punsly, Brian; Multi-Epoch Observations of the Redwing Excess in the Spectrum of 3C279 2013 ApJ Letters 762 25

Punsly, B. and Rodriguez, J. The Relationship Between X-ray Luminosity and Major Flare Launching in GRS~1915+105 2013 ApJ 764 173

Punsly, B. and Rodriguez, J. GRS 1915+105 as a Galactic Analog of a Fanaroff-Riley II Quasar ApJ 770 90

Reynolds, Cormac; Punsly, Brian; O'Dea, Christopher P. Misalignment of the Jet and the Normal to the Dusty Torus in the Broad Absorption Line QSO FIRST J155633.8+351758 ApJL 773 10

Reynolds, Cormac; Punsly, Brian; O'Dea, Christopher P.; Hurley-Walker, Natasha A Blazar-like Radio Flare in Mrk 231 ApJL 776 21

Punsly, B. and Rodriguez, J. Calibrated estimates of the energy in major flares of GRS 1915+105 MNRAS 435 2322